

Bacteria TMDL in the Piney Run Watershed

First Public Meeting December 18, 2003



Presentation Overview

- 1. Overview of Virginia's TMDL Program
- 2. Applicable Water Quality Standard
- 3. Piney Run Impairment
- 4. TMDL Development Approach
- 5. Bacteria Source Assessment

What is a TMDL?

- TMDL stands for **Total Maximum Daily Load**
- A TMDL is a pollution budget
- A TMDL is a calculation of the maximum amount of a pollutant that a waterbody can receive and still meet water quality standards
- A TMDL includes an **allocation** of that maximum amount to the pollutant's sources

TMDL Equation

A TMDL is summarized as:

TMDL = Sum of WLA + Sum of LA + MOS

Where:

- TMDL = Total Maximum Daily Load
- WLA = Waste Load Allocation (point sources)
- Load Allocation (nonpoint sources)
- -MOS = Margin of Safety

How is a TMDL developed?

- Identify all sources of a given pollutant within the watershed
- Calculate the amount of pollutant entering the stream from each source
- Calculate the pollutant reductions needed, by source, to attain water quality standards
- Allocate the allowable loading to each source and include a margin of safety

When are TMDLs needed?

- State and federal law require TMDLs to be developed for **impaired** waters
- Impaired waters do not meet applicable water quality standards (WQS)
- Waters that do not meet WQS do not support their **designated use(s)**
- For bacteria impairments, the designated use that is affected is the **recreational use**

Regulatory Basis of TMDLs

- TMDLs required by Federal and State law
 - 1972 Clean Water Act (CWA), Section 303(d)
 - 1997 Water Quality Monitoring, Information and Restoration Act (WQMIRA)
- 1998 lawsuit filed by the American Canoe Association and the American Littoral Society against EPA for failure to comply with CWA §303(d) in Virginia
- 1999 Consent Decree requiring EPA and Virginia to complete 636 TMDLs by 2010

Regulatory Requirements

- Both state and federal law require:
 - Establishment of water quality standards
 - Monitoring of water quality in surface waters
 - Assessment of water quality in surface waters
 - Listing of waters that do not meet water quality standards (impaired waters)
 - Development of TMDLs for impaired waters
- State law requires, and federal law recommends:
 - Development of a TMDL Implementation Plan

Roles of DEQ and DCR in TMDL and IP Development

- DEQ is the lead for TMDL development, including submittal to EPA
- DCR is the lead for TMDL Implementation Plan (IP) development
- DEQ is responsible for ensuring public participation in the TMDL program

Presentation Overview

- 1. Overview of Virginia's TMDL Program
- 2. Applicable Water Quality Standard
- 3. Piney Run Impairment
- 4. TMDL Development Approach
- 5. Bacteria Source Assessment

Water Quality Standards

- Water Quality Standards (WQS):
 - set by states and approved by EPA
 - set numeric and narrative limits on pollutants
 - consist of designated use(s) and water quality
 criteria
- Purpose of WQS:
 - protection of 5 designated uses (aquatic life, fish consumption, shellfish, recreation, drinking water)
 - restoration of state waters to meet criteria

Applicable Designated Use


- All surface waters in Virginia are currently designated for primary contact recreation (e.g. swimming)
- In March 2003, a **secondary contact recreation** use designation (e.g. wading, fishing) was added to the WQS
 - Five times the primary contact criteria
 - Individual waters will only be considered for reclassification after TMDL implementation has been tried using reasonable BMPs
 - Effective date pending EPA approval

Pollutant of Concern

- *Fecal bacteria* are found in the digestive tract of humans and warm blooded animals
- Fecal bacteria are an indicator of the potential presence of pathogens in waterbodies
- The presence of fecal bacteria in water samples is a strong indicator of recent sewage or animal waste contamination

Sampling for Bacteria

- Stream samples are collected in sterile 125 mL sample bottles
- Samples are filtered to deposit bacteria on filters
- Filters are incubated, allowing individual bacteria to grow into visible colonies
- Colonies are counted to give a concentration of colony forming units (cfu) per 100 mL



Old Criteria

- Indicator species: fecal coliform
 - used in listing Piney Run
- Instantaneous max: 1,000 cfu/100 mL
- Applicable for data sets with 1 or fewer samples in 30 days
- Geometric mean: 200 cfu/100 mL
- Applicable for data sets with 2 or more samples in 30 days

New Criteria

- Indicator species for freshwater: *E. coli*
 - change in indicator species from fecal coliform to *E. coli* (fresh water)
 - E. coli bacteria are a subset of fecal coliform bacteria and correlate better with swimming-associated illness
- Instantaneous max: 235 cfu/100 mL
- Applicable for all data sets; no samples may exceed the maximum
- Geometric mean: 126 cfu/100 mL
- Applicable for data sets with 2 or more samples in a calendar month

Summary of Changes in Primary Contact Criteria

| Indicator | Status | Instantaneous Maximum (cfu/100mL) | Geometric Mean (cfu/100 mL) |
|----------------|---------|---|-----------------------------------|
| Fecal Coliform | Old | 1,000 | 200 |
| E. coli | New | 235 | 126 |
| Fecal Coliform | Interim | 400 | 200 |

- Changes went into effect on January 15, 2003
- Both New *E. coli* and Interim Fecal Coliform criteria apply
- Fecal coliform criteria will be phased out entirely once 12 *E. coli* samples have been collected or after June 30, 2008

Comparison of the Old Fecal Coliform and New *E. coli* Criteria

| Old FC (cfu/100mL) | Interim FC (cfu/100mL) | FC translated to EC* (cfu/100mL) | New EC (cfu/100mL) |
|-----------------------|---------------------------|--|-----------------------|
| 200 | 200 | 129 | 126 |
| | 400 | 243 | 235 |
| 1,000 | | 565 | |

^{*} Based on regression model between 493 dual data points

Note: FC = Fecal Coliform, EC = *Escherichia Coli*


Presentation Overview

- 1. Overview of Virginia's TMDL Program
- 2. Applicable Water Quality Standard
- 3. Piney Run Impairment
- 4. TMDL Development Approach
- 5. Bacteria Source Assessment

Impairment in the Piney Run Watershed


| WATER | CAUSE | STREAM | LENGTH | YEARS |
|----------|----------|---|---------|------------|
| BODY | | NAME | (Miles) | LISTED |
| VAN-A01R | Bacteria | Piney Run (from mouth of unnamed lake to confluence with Potomac River) | 3.52 | 1998, 2002 |

Map of the Piney Run Watershed



- DEQ monitoring station: 1APIA001.80
- USGS flow gage: 01636690
- 2002 305(b) results: 5 of 22 samples (23%) exceeding 1000 cfu/100mL
- 2000 305(b) results: 5 of 20 (25%)
- 1998 305(b) results: 5 of 19 (26%)

Fecal Coliform in Piney Run (1APIA001.80)



Presentation Overview

- 1. Overview of Virginia's TMDL Program
- 2. Applicable Water Quality Standard
- 3. Piney Run Impairment
- 4. TMDL Development Approach
- 5. Bacteria Source Assessment

What is Load Duration Analysis?

- Less complex spreadsheet model for TMDL development
- Approach proposed for bacteria TMDLs in small watersheds
- Model requires
 - stream flow data
 - ambient water quality data, and
 - bacteria source tracking data (for pollutant source identification and loading allocations)


Development of Flow Duration Curve for Piney Run

- Piney Run has a USGS flow gaging station that was established in 2001
- In order to include the time period that led to the listing (1/1/1996 to 12/31/2000 for the most recent assessment), the flow record must be extended

Reference Stream Selection

- Flows were correlated with Catoctin, Goose and Passage Creeks
- The period from 1988 to present was used
- Piney Run flows correlated best with Catoctin Creek (0.9317)
- Flow regression equations were then used to generate continuous flow records (1988-03)


Piney Run Flow Duration Curve



Load Duration Curve

- Represents the maximum amount of a pollutant allowed at each flow level
- Obtained by multiplying the flow duration curve by the water quality criterion
- At higher flows, a stream will have more assimilative capacity
- At lower flows, it will have less assimilative capacity


Piney Run Load Duration Curve




TMDL Required Reduction

- The TMDL must ensure water quality is protected during times when stream is most vulnerable
- The stream is assumed to be most vulnerable when the highest exceedance occurs
- This critical condition occurred on 9/26/2000, with an observed concentration of 3,819 cfu/100mL at an estimated flow of 25.72 cfs

Piney Run Load Duration Curve



Piney Run TMDL



TMDL Reduction Required

- The TMDL equation is then calculated using the maximum observed exceedance and average flow conditions (10.23 cfs)
- At average flow:

```
- the existing load is 3.49 \times 10^{14} \text{ cfu/yr}
```

- the allowable load is $2.15 \times 10^{13} \text{ cfu/yr}$
- the required reduction is $3.28 \times 10^{14} \text{ cfu/yr}$
- This corresponds to a 94% reduction


Development of TMDL Allocations

- Assume an implicit margin of safety due to conservative assumptions
- Subtract point source loads from the TMDL load to obtain the non-point source load
- Use results of source assessment and BST study to allocate the non-point source loads among sources (human, livestock, wildlife)

Presentation Overview

- 1. Overview of Virginia's TMDL Program
- 2. Applicable Water Quality Standard
- 3. Piney Run Impairment
- 4. TMDL Development Approach
- 5. Bacteria Source Assessment

Land Use in the Piney Run Watershed



| MRLC | Piney | Run | |
|------------------|-------|---------|--|
| Land Use | Acres | Percent | |
| Cropland | 48 | 0.5% | |
| Pasture | 2,616 | 26.9% | |
| Barren or Mining | 0 | 0.0% | |
| Forest | 6,908 | 71.0% | |
| Transitional | 1 | 0.0% | |
| Urban | 20 | 0.2% | |
| Water | 36 | 0.4% | |
| Wetlands | 104 | 1.1% | |
| Total | 9,731 | 100.0% | |

Potential Sources of Bacteria in Piney Run

- Humans/Pets
 - Straight Pipes
 - Septic Systems
 - Biosolids
 - Permitted PointSources
 - Pets

- Livestock
 - Direct Deposit to Land and Streams
 - Land Application
- Wildlife
 - Direct Deposit to Land and Streams

Potential Human and Pet Sources



Estimated Human and Pet Sources

| Source | Piney Run | Reference |
|----------------|-----------|--------------------------|
| People | 626 | 2000 Census |
| Dogs | 411 | 2000 Census, APPMA |
| VPDES Permits | 0 | VADEQ |
| SFH Permits | 2 | VADEQ |
| Septic Systems | TBD | Loudoun Co. Health Dept. |
| Straight Pipes | TBD | Loudoun Co. Health Dept. |
| Biosolids | TBD | Loudoun Co. Health Dept. |

| Watershed | Permit No. | Facility Name | Design Flow |
|-----------|------------|--------------------------------------|--------------------|
| Piney Run | VAG406106 | Neersville Volunteer Fire and Rescue | 400 gal/day |
| Piney Run | VA G406249 | Amoco - Tri State | 875 gal/day |

Potential Livestock Sources



Estimated Livestock Sources

| Source | Piney Run | Reference |
|-------------------|-----------|-----------------------------|
| Cattle and calves | 500 | 1997 Ag Census/Loudoun SWCD |
| Beef Cows | 225 | 1997 Ag Census/Loudoun SWCD |
| Hogs and Pigs | 0 | 1997 Ag Census/Loudoun SWCD |
| Sheep and Lambs | 30 | 1997 Ag Census/Loudoun SWCD |
| Layers | 50 | 1997 Ag Census/Loudoun SWCD |
| Broilers | 0 | 1997 Ag Census/Loudoun SWCD |
| Horses | 350 | Loudoun SWCD |

• Livestock numbers estimates based on discussion with the Loudoun Soil and Water Conservation District (12/11/2003)





Estimated Wildlife Sources

| Animal | Habitat | Density | Piney Run |
|----------|--------------------------|----------------|-----------|
| Deer | Forest, Agriculture, | 0.084 per acre | 806 |
| | Urban Pervious | | |
| Raccoons | Within 600 ft of streams | 0.07 per acre | 119 |
| Muskrats | Within 66 ft of streams | 2.75 per acre | 531 |
| Beavers | Streams | 4.8 per mile | 66 |
| Turkeys | Forest | 0.01 per acre | 69 |
| Ducks | Within 66 ft of streams | 0.008 per acre | 2 |
| Geese | Within 66 ft of streams | 0.02 per acre | 4 |

• Wildlife numbers estimated based on habitat types and animal densities from the Catoctin and Goose Creek bacteria TMDLs

Bacteria TMDL for the Piney Run Watershed

- First public meeting:
 - Thursday, December 18
 - Neersville Fire and Rescue Building, 11762 Harpers Ferry Road, Hillsboro, Virginia
 - First 30 day comment period ends January 16, 2004
- Second and final public meeting will be held in February 2004
 - Results of BST study
 - Draft report (Second 30 day comment period)
- Submit to EPA for approval

Bacteria TMDL for the Piney Run Watershed

Kate Bennett
Regional TMDL Coordinator
Northern Virginia Regional Office
VA Department of Environmental Quality
13901 Crown Ct.

Woodbridge, VA 22193

Phone: (703) 583-3896

Fax: (703) 583-3841

E-mail: kebennett@deq.state.va.us

Interim Criteria

- Indicator species: fecal coliform
 - will be phased out when 12 *E. coli* observations available or after June 30, 2008, whichever comes first
 - will not be used to assess compliance
- Instantaneous max: 400 cfu/100 mL
- Applicable for all data sets; no more than 10% of samples in a calendar month may exceed the maximum
- Geometric mean: 200 cfu/100 mL
 - Applicable for data sets with 2 or more samples in a calendar month